

Three new species of the leafhopper genus *Arboridia* Zachvatkin (Hemiptera, Cicadellidae, Typhlocybinae), with a key and checklist to known species of China

Chang Han¹, Bin Yan¹, Xiaofei Yu², Maofa Yang^{1,2}, Michael D. Webb³

¹ Institute of Entomology, Guizhou Provincial Key Laboratory for Agricultural Pest Management of the Mountainous Region, Guizhou University, Guiyang 550025, China

² College of Tobacco Science, Guizhou University, Guiyang 550025, China

³ Department of Entomology, The Natural History Museum, Cromwell Road, London, SW7 5BD, UK

Corresponding authors: Maofa Yang (gddgly@126.com); Michael D. Webb (M.Webb@nhm.ac.uk)

Abstract

Three new species of the leafhopper genus *Arboridia* Zachvatkin 1946, *Arboridia* (*Arboridia*) *furcata* Han, **sp. nov.**, *Arboridia* (*Arboridia*) *rubrovittata* Han, **sp. nov.**, and *Arboridia* (*Arboridia*) *robustipenis* Han, **sp. nov.**, are described and illustrated from fruit trees in Southwest China. A key and checklist to known species from China are provided.

Key words: *Arboridia*, Hemiptera, identification key, new species, taxonomy, Typhlocybinae



Academic editor: Ilia Gjonov

Received: 15 January 2024

Accepted: 27 February 2024

Published: 28 March 2024

ZooBank: <https://zoobank.org/91F2B2E0-AC9E-4FA1-8487-BFB3BF86CF21>

Citation: Han C, Yan B, Yu X, Yang M, Webb MD (2024) Three new species of the leafhopper genus *Arboridia* Zachvatkin (Hemiptera, Cicadellidae, Typhlocybinae), with a key and checklist to known species of China. ZooKeys 1196: 255–269. <https://doi.org/10.3897/zookeys.1196.118829>

Copyright: © Chang Han et al.

This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Introduction

The leafhopper genus *Arboridia* Zachvatkin, 1946 belongs to the tribe Erythroneurini of the subfamily Typhlocybinae (Hemiptera: Auchenorrhyncha: Cicadellidae) and includes two subgenera, *Arboridia* Zachvatkin, 1946 and *Arborifera* Sohi & Sandhu, 1971. Species feed on a variety of plants including fruit trees, hawthorn, maple, honeysuckle, dogwood and several other plants (Song and Li 2013). So far, 84 species have been described in this large genus, distributed throughout the Palaearctic and Oriental regions, including 25 valid species from China (Song et al. 2016; Cao et al. 2019).

In this study, three new species are described from Guizhou, China. Photographs of the whole body, illustrations of male genitalia, and biological information such as host plants and distributional records are provided. In addition, an updated key to the *Arboridia* species from China is provided.

Materials and methods

Specimens used in this study were collected from grape, kiwi and walnut trees in Guizhou, China using a sweep net. A Nikon SMZ 1270 microscope was used to dissect the specimens and an Olympus CX41 microscope for observing and drawing the male genitalia. A KEYENCE VHX-6000 digital microscope was used to take pictures of the male habitus. Morphological terminology used in this

work follows Dietrich (2005) and Dworakowska (1993). All specimens examined in this study are deposited in the Institute of Entomology, Guizhou University, China (GUGC).

Taxonomy

Arboridia Zachvatkin

Arboridia Zachvatkin, 1946: 153. Type species. *Typhlocyba parvula* Boheman, 1845. *Khoduma* Dworakowska, 1972: 403. Synonymised by Dworakowska and Virak-tamath (1975: 529). Type species. *Khoduma jacobii* Dworakowska, 1972.

Diagnosis. Head slightly narrower than pronotum, crown weakly produced with fore margin rounded. Head and thorax yellow; vertex usually with pair of dark subapical spots; pronotum usually with irregular brown symmetric markings; scutellum with brown basal triangles. Forewing either without marking, with oblique vittae or with dark spots. Ventral abdominal apodemes small and extended to or beyond posterior margin of 3rd sternite. Male pygofer with wide-spread microtrichia and several small rigid setae on inner surface of hind margin; dorsal appendage present, free from pygofer side; ventral appendage absent; phragma lobe with spine-like setae present on each side of aedeagus, attached to dorsal apodeme of aedeagus by ligaments (Fig. 51). Subgenital plate upturned apically with lateral margin basally expanded triangular shaped with 2–4 lateral macrosetae in an oblique row slightly basad of midlength; lateral margin with short spine-like setae. Style apex usually with 3 points, sometimes 2nd point absent. Aedeagus with shaft laterally compressed, usually with processes, gonopore apical on ventral surface; dorsal apodeme and preatrium present or absent. Connective U- or V- shaped with median anterior lobe absent.

Distribution. Palaearctic and Oriental regions.

Checklist of Chinese species of *Arboridia*

1. *Arboridia agrillacea* (Anufriev, 1969b: 182–183, fig. 13: 1–6, *Erythroneura*); Anufriev, 1978a: 87, transferred to *Arboridia*; Song & Li, 2013: 243–244, figs J, j, jj, 63–69; *Arboridia koreana* Oh & Jung, 2015: 447–448, figs 1, 3, 5, 7, 9–15, synonym. Distribution: Gansu, Guangxi, Guizhou, Henan, Shaanxi, Sichuan.
2. *Arboridia anteoculara* Song & Li, 2013: 230–233, figs A, a, 1–7. Distribution: Guizhou.
3. *Arboridia apicalis* (Nawa, 1913a: 480–486, Pl. 24, *Zygina*); Cockerell, 1920a: 247, *Erythroneura*; *Erythroneura sandagouensis* Vilbaste, 1968a: 108, synonym; Anufriev, 1969b: 185–186, fig. 15: 8–13; Dworakowska, 1970g: 607–608, fig. 18, transferred to *Arboridia*. Distribution: Anhui, Guizhou, Hebei, Henan, Hubei, Jiangsu, Liaoning, Shannxi, Shandong, Taiwan, Zhejiang.
4. *Arboridia baiyunensis* Song & Li, 2013: 233–234, figs B, b, 8–14. Distribution: Henan.
5. *Arboridia (Arborifera) changlingensis* Jiang, Luo & Song, 2021: 354–355, figs 5–8, 27–34. Distribution: Guizhou.

6. *Arboridia cincta* Song & Li, 2015: 585–587, figs A–C, 1–7. Distribution: Henan.
7. *Arboridia cuihuashana* Song & Li, 2013: 237–238, figs E, e, 29–35. Distribution: Shaanxi.
8. *Arboridia echinata* Song & Li, 2013: 239–240, figs G, g, gg, 42–48. Distribution: Guizhou.
9. *Arboridia furcata* Han, sp. nov. Distribution: Guizhou.
10. *Arboridia huajiangensis* Jiang, Luo & Song, 2021: 351–353, figs 1–4, 9–26. Distribution: Guizhou.
11. *Arboridia jinghongensis* Pu, Wang & Song, 2023: 296–297, figs 1a–f, 2a–h. Distribution: Yunnan.
12. *Arboridia kakogawana* (Matsumura, 1932: 113, *Zygina*); Ishihara, 1953b: 33, *Erythroneura*; Dworakowska, 1970g: 610, figs 25–29, transferred to *Arboridia*. Distribution: Beijing, Guizhou, Shandong, Xinjiang.
13. *Arboridia lunula* Song & Li, 2013: 234–236, figs D, d, 22–28. Distribution: Guizhou.
14. *Arboridia luojiashangensis* Zhang, Jiang & Song, 2022: 6–8, figs 21–32. Distribution: Guizhou.
15. *Arboridia maculifrons* (Vilbaste, 1968a: 107, *Erythroneura*); Dworakowska, 1970g: 611, figs 19–22, transferred to *Arboridia*. Distribution: Guizhou, Hebei.
16. *Arboridia ochracea* Song & Li, 2015: 587–588, figs D–F, 8–15. Distribution: Henan.
17. *Arboridia paraprocesa* Song & Li, 2013: 239, figs F, f, 36–41. Distribution: Guizhou, Henan.
18. *Arboridia reniformis* Song & Li, 2013: 234, figs C, c, cc, 15–21. Distribution: Yunnan.
19. *Arboridia remmi* (Vilbaste, 1968a: 103, *Erythroneura*); Anufriev, 1969b: 183–184, figs 15: 1–7; Dworakowska, 1970g: 613, transferred to *Arboridia*. Distribution: Guizhou.
20. *Arboridia robustipenis* Han, sp. nov. Distribution: Guizhou.
21. *Arboridia rubrovittata* Han, sp. nov. Distribution: Guizhou.
22. *Arboridia sinensis* Guglielmino, Xu, Buckle & Dong, 2012: 878–881, figs 1: A–F, 2: A–B. Distribution: Yunnan.
23. *Arboridia suputinkaensis* (Vilbaste, 1968a: 109, *Erythroneura*); Dworakowska, 1970g: 613, transferred to *Arboridia*. Distribution: Henan, Zhejiang. <https://hoppers.speciesfile.org/otus/43920/overview> (Dmitriev et al. 2022)
24. *Arboridia* (*Arborifera*) *surstyli* Cai & Xu, 2006: 75–76, figs 1: 1–10. Distribution: Henan, Zhejiang.
25. *Arboridia suzukii* (Matsumura, 1916b: 396, *Zygina*); Ishihara, 1953b: 34, *Erythroneura*; *Erythroneura arboricola* Vilbaste, 1968a: 101, synonym; Dworakowska, 1970g: 613, transferred to *Arboridia*. Distribution: Gansu, Guizhou, shannxi, shanxi, Taiwan. <https://hoppers.speciesfile.org/otus/43922/overview>.
26. *Arboridia tridentata* Song & Li, 2013: 240–241, figs H, h, 49–55. Distribution: Yunnan.
27. *Arboridia xiaotungensis* Zhang, Jiang & Song, 2022: 2–5, figs 1–20. Distribution: Guizhou.
28. *Arboridia zhenyuana* Song & Li, 2013: 242–243, figs I, i, 56–62. Distribution: Gansu.

Key to species (males) of *Arboridia* species from China

(modified from Jiang et al. 2021)

- 1 Preatrium of aedeagus short or absent (*Arborifera*)2
- Preatrium of aedeagus well developed (Figs 24, 39, 55) (*Arboridia*)3
- 2 Aedeagal shaft with pair of sharp inverted processes on dorsal margin.....*A. surstyli*
- Aedeagal shaft with one broad triangular process on dorsal margin.....*A. changlingensis*
- 3 Aedeagus without process, shaft with pair of lateral flanges4
- Aedeagus with processes, shaft without pair of lateral flanges6
- 4 Aedeagal shaft with lateral flanges serrate *A. zhenyuana*
- Aedeagal shaft with lateral flanges not serrate.....5
- 5 Aedeagal shaft with lateral flanges narrow, entire*A. agrillacea*
- Aedeagal shaft with larger lateral flanges partly wrapped around shaft.....*A. jinghongensis*
- 6 Aedeagus with one process7
- Aedeagus with more than one process9
- 7 Aedeagus with process arising from preatrium *A. apicalis*
- Aedeagus with process arising from midlength of shaft.....8
- 8 Aedeagus with dorsal apodeme extremely expanded in lateral view*A. sinensis*
- Aedeagus with dorsal apodeme narrow in lateral view*A. tridentata*
- 9 Aedeagus with one or two pairs of processes10
- Aedeagus with three or more processes19
- 10 Aedeagal shaft with two pairs of processes, one at apex and one at base ..*A. ochracea*
- Aedeagal shaft with one pair of processes arising from apex or base11
- 11 Aedeagus with processes arising from base of shaft12
- Aedeagus with processes arising from apex of shaft15
- 12 Aedeagus with two pairs of basal processes.....*A. anteoculara*
- Aedeagus with one pair of basal processes.....13
- 13 Aedeagus with processes slender and bent basad apically (Figs 17, 24).....*A. furcata* sp. nov.
- Aedeagus with processes stout and straight14
- 14 Aedeagus with dorsal apodeme narrow in lateral view*A. lunula*
- Aedeagus with dorsal apodeme extremely expanded in lateral view*A. maculifrons*
- 15 Aedeagus with apical processes directed basally16
- Aedeagus with apical processes directed distally18
- 16 Apex of aedeagal shaft acuminate in ventral view..... *A. cincta*
- Apex of aedeagal shaft truncate in ventral view17
- 17 Aedeagus without subapical bifurcation in ventral view*A. reniformis*
- Aedeagus with subapical bifurcation in ventral view.....*A. xiaotungensis*
- 18 Aedeagal shaft without spines*A. cuihuashana*
- Aedeagal shaft with numerous short spines.....*A. echinata*
- 19 Aedeagal shaft with two or three processes at midlength.....20
- Aedeagal shaft with one pair of apical processes22

- 20 Aedeagal shaft with three processes subbasally, a pair of upper bifurcate processes and a slightly more ventral process (Figs 35, 40) ***A. rubrovittata* sp. nov.**
- Aedeagal shaft with two processes at midlength **21**
- 21 Aedeagal shaft with two processes fused for 2/3 of their length (Figs 50–51, 55–56) ***A. robustipenis* sp. nov.**
- Aedeagal shaft with two processes one above the other... ***A. luojiashangensis***
- 22 Apical processes of aedeagal shaft directed basally **23**
- Apical processes of aedeagal shaft directed distally **24**
- 23 Aedeagal shaft with slender apical processes, without ventral protrusion...
..... ***A. suputinkaensis***
- Aedeagal shaft with short apical processes, with small ventral protrusion..
..... ***A. huajiagensis***
- 24 Preatrium of aedeagus longer than shaft **25**
- Preatrium of aedeagus shorter than shaft **26**
- 25 Aedeagal shaft with numerous short spines ***A. suzukii***
- Aedeagal shaft without spines ***A. remmi***
- 26 Aedeagal shaft with distinct extension at midlength ***A. baiyunensis***
- Aedeagal shaft without extension at midlength **27**
- 27 Aedeagus with dorsal apodeme and shaft narrow in lateral view; preatrium with a long ventral process ***A. paraprocesa***
- Aedeagus with dorsal apodeme and shaft expanded in lateral view; preatrium with a short ventral process ***A. kakogawana***

***Arboridia (Arboridia) furcata* Han, sp. nov.**

<https://zoobank.org/92B51DA2-4F3F-40E6-9F01-D8D34BAD4884>


Figs 1–4, 14–28

Description. Dorsum yellowish brown; eyes grey with posterior margin beige; vertex with a pair of black spots subapically; coronal suture indistinct distally, pale brown basally (Figs 1–3). Face yellowish brown with median area of frontoclypeus and anteclypeus brighter towards apex; lorum and gena whitish (Fig. 4). Pronotum yellowish brown with brownish spots at anterior margin. Scutellum yellow with lateral triangles dark brown (Fig. 3). Forewing hyaline, veins brown. Abdominal tergites black; sternites milky white; subgenital plate dark apically (Figs 2, 14).

Ventral abdominal apodemes small, extended to 4th sternite (Figs 18, 27).

Male genitalia. Pygofer dorsal appendage simple, slender and wavy, with the apex obliquely truncate (Figs 14, 21). Subgenital plate with 3 lateral macrosetae in an oblique row slightly basad of midlength laterally (Figs 15, 16, 22, 23). Style long and slender, apex with 3 points; preapical lobe well developed; several small tubercles subapically and at midlength (Figs 20, 26). Aedeagal shaft long and stout, slightly laterally compressed, a pair of long slender basal processes on ventral surface of the shaft, parallel to the shaft in their basal half, then sharply turned in proximal direction in their distal half (Figs 17, 19, 24, 25); dorsal apodeme short and robust, expanded laterally at apex; preatrium short (Figs 17, 24). Connective U-shaped, with lateral arms long and stem broad (Figs 19, 28).

Measurement. Body length males 3.0–3.2 mm, females 3.2–3.3 mm.



Figures 1–13. External morphology of *Arboridia* species 1–4 *Arboridia furcata* Han, sp. nov. 1 habitus, dorsal view 2 habitus, lateral view 3 head and thorax, dorsal view 4 face 5–9 *Arboridia rubrovittata* Han, sp. nov. 5 habitus, dorsal view 6 habitus, lateral view 7, 8 head and thorax, dorsal view 9 face 10–13 *Arboridia robustipenis* Han, sp. nov. 10 habitus, dorsal view 11 habitus, lateral view 12 head and thorax, dorsal view 13 face. Scale bars: 0.5 mm.




Figures 14–20. *Arboridia* (A.) *furcata* Han, sp. nov. **14** male genitalia, lateral view **15** subgenital plate, dorsal view **16** subgenital plate, ventral view **17** aedeagus, connective and style, lateral view **18** abdominal apodemes **19** aedeagus, connective and style, ventral view **20** style, lateral view. Scale bars: 0.1 mm.

Specimen examined. Holotype ♂: CHINA, Guizhou Prov., Dejiang, 22.VII.2017, coll. Chang Han and Bin Yan, on grape (GUGC). **Paratypes:** 5♂♂, 5♀♀, same data as holotype.

Etymology. The new species is named from the Latin word “*furcatus*”, referring to the forked aedeagal process.

Remarks. The new species is similar to *Arboridia* (A.) *anteoculara* Song & Li, 2013, but differs in only having a pair of processes on the ventrobasal surface of aedeagal shaft (Figs 17, 24); the latter species having two pairs of processes and arising from both sides of the aedeagal shaft.

Host. *Vitis vinifera* L. (grape).



Figures 21–28. *Arboridia* (*A.*) *furcata* Han, sp. nov. **21** male pygofer, lateral view **22** subgenital plate, ventral view **23** subgenital plate, dorsal view **24** aedeagus, lateral view **25** aedeagus, ventral view **26** style, lateral view **27** abdominal apodemes **28** connective.

***Arboridia (Arboridia) rubrovittata* Han, sp. nov.**


<https://zoobank.org/AE69B8DF-EB76-4CB8-A0EA-B9EB4D1FCB43>

Figs 5–9, 29–44


Description. Dorsum orange, yellow or beige. Eyes black, inner and posterior margins white (Figs 5, 6). Vertex without pair of dark spots, with a white patch each side of midline basally; coronal suture orange yellow (Figs 5, 7, 8). Face orange yellow. Pronotum with ivory or pale white streaks situated laterad of anterior margin.



Figures 29–36. *Arboridia (Arboridia) rubrovittata* Han, sp. nov. **29** male genitalia, lateral view **30** style, lateral view **31** subgenital plate **32** abdominal apodemes **33** aedeagus, connective and style, lateral view **34** aedeagus, lateral view **35** aedeagus, ventral view **36** aedeagus, connective and style, ventral view. Scale bars: 0.1 mm.



Figures 37–44. *Arboridia (Arboridia) rubrovittata* Han, sp. nov. **37** male pygofer, lateral view **38** subgenital plate **39** aedeagus, lateral view **40** aedeagus, ventral view **41** style, lateral view **42** style apex, ventral view **43** abdominal apodemes **44** connective.



Figures 45–52. *Arboridia (Arboridia) robustipenis* Han, sp. nov. **45** male genitalia, lateral view **46** style, lateral view **47** subgenital plate **48** aedeagus, connective and style, lateral view **49** abdominal apodemes **50** aedeagus, lateral view **51** aedeagus & pygofer dorsal appendage, ventral view **52** aedeagus, connective and style, ventral view. Scale bars: 0.1 mm.

Scutellum pale or whitish yellow with lateral triangles dark to pale brown (Figs 5, 7, 8). Forewing with oblique pale reddish-orange vittae in clavus and adjacent area of wing. Abdominal segments milky yellow, tergites with segment margins brown. Subgenital plate with nearly 2/3 apically dark (Figs 5, 6).

Ventral abdominal apodemes small, extended to posterior margin of 3rd sternite (Figs 32, 43).

Male genitalia. Pygofer dorsal appendage claw-like (Figs 29, 37). Subgenital plate with 3 lateral macrosetae in an oblique row slightly basad of midlength

(Figs 31, 38). Style slender, with 2 points, heel point small; sword-like apically with transverse wrinkles in lateral view (Figs 30, 41), serrated in ventral view (Fig. 42). Aedeagus relatively small, shaft laterally compressed, digitate and slightly upturned in lateral view; subbasally with three processes, two basal processes and a single unpaired spike basad, the distal paired processes divergent with branches slender, the proximal process slightly shorter and more robust, finger-like in ventral view (Figs 34, 35, 39, 40); preatrium long. Connective V-shaped (Figs 36, 44).

Measurement. Body length males 2.7–2.9 mm, females 2.9–3.0 mm.

Specimen examined. *Holotype* ♂: CHINA, Guizhou Prov., Jianhe, 26.V.2017, coll. Chang Han and Yaowen Zhang, on kiwi (GUGC). *Paratypes*: 23♂♂, 25♀♀, same data as holotype on kiwi; 3♂♂, 6♀♀, CHINA, Guizhou Prov., Wujiang, 19.V.2017, coll. Chang Han and Bin Yan, on walnut (GUGC).

Etymology. The new species name is derived from the Latin words “*ruber*” (red) and “*vittatus*” (banded), referring to the reddish-orange oblique stripes on the forewings.

Remarks. The new species can be distinguished from most *Arboridia* species by its vertex and pronotum without dark spots (Figs 5, 7, 8) and reddish-orange stripes on the forewing. Its male genitalia is similar to *A. (A.) lunula* Song & Li, 2013, but can be distinguished by the sword-like apex of the style and aedeagus with three basal ventral processes, the upper paired processes slender (Figs 34, 39).

Host. *Actinidia chinensis* Planch (kiwi fruit); *Juglans regia* L. (walnut).

***Arboridia (Arboridia) robustipenis* Han, sp. nov.**

<https://zoobank.org/C2F0CB93-2BA0-498A-878F-F91A7FE2D8DD>


Figs 10–13, 45–59

Description. Head with eyes black with posterior margin pearl white; crown yellow with a dark yellow spot at apex, an adjacent brown spot posteriorly on each side of midline and a brown patch at base of coronal suture (Figs 10–12). Face pale yellow, with anteclypeus brown apically; gena whitish yellow (Fig. 13). Pronotum semitransparent with brown markings (Fig. 12). Scutellum yellow with lateral triangles dark brown (Fig. 12). Forewing brownish hyaline with off-white patch in clavus and brochosome region. Abdominal segments dark brown, sternites with yellow hind margins of segments; subgenital plates black apically (Figs 10, 11).

Abdominal apodemes small, not exceeding posterior margin of 3rd sternite (Figs 49, 58).

Male genitalia. Pygofer dorsal appendage tapering and curved ventrad (Figs 45, 53). Subgenital plate with 3 lateral macrosetae in an oblique row slightly basad of midlength (Figs 47, 54). Style apex with two triangular points; preapical lobe well developed (Figs 46, 57). Aedeagal shaft strongly laterally compressed and “C” shaped with apex upturned in lateral view, with two basal processes fused for 2/3 of their length at midlength of ventral margin; preatrium long, with a thorn-like basal projection (Figs 50–51, 55–56). Connective V-shaped with stem reduced (Figs 52, 59).

Measurement. Body length males 2.9–3.2 mm, females 3.0–3.3 mm.



Figures 53–59. *Arboridia (Arboridia) robustipenis* Han, sp. nov. **53** male pygofer, lateral view **54** subgenital plate **55** aedeagus, lateral view **56** aedeagus, ventral view **57** style, lateral view **58** abdominal apodemes **59** connective.

Specimen examined. Holotype: ♂, CHINA, Guizhou Prov., Wujiang, 19.V.2017, coll. Chang Han and Bin Yan, on walnut (GUGC). **Paratypes:** 5♂♂, 7♀♀, same data as holotype on walnut; 50♂♂61♀♀, CHINA, Guizhou Prov., Xiuwen, 19.VII.2017, coll. Chang Han and Bin Yan, on kiwi (GUGC)

Etymology. The new species name is derived from the Latin words “robustus” and “penis”, and refers to the robust aedeagal shaft in lateral view.

Remarks. The new species can be distinguished from *A. (A.) luojiashangensis* Zhang & Song, 2022 by the aedeagus with strongly laterally compressed shaft “C” shaped; the paired basal processes fused for 2/3 of their length like a forked tongue (Figs 50–51, 55–56); and the preatrium with a thorn-like basal projection.

Host. *Actinidia chinensis* Planch (kiwi fruit); *Juglans regia* L. (walnut).

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

The study was supported by the Program of Natural Science Foundation of China (No. 32360393, 31802002), Guizhou Province Science and Technology Innovation Talent Team Project (Qian Ke He Pingtai Rencai-CXTD [2021]-004) and the Basic Research Program (the natural science projects) of Guizhou Province (ZK[2022]-125).

Author contributions

Conceptualization: XY, CH. Data curation: CH. Formal analysis: CH. Funding acquisition: XY, MY. Investigation: CH, BY. Methodology: CH. Project administration: CH. Writing - original draft: CH. Writing - review and editing: MW, XY, MY.

Author ORCIDs

Bin Yan  <https://orcid.org/0000-0002-5290-4576>

Maofa Yang  <https://orcid.org/0000-0002-5523-6825>

Michael D. Webb  <https://orcid.org/0000-0000-0000-0000>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Cao YH, Dmitriev DA, Dietrich CH, Zhang YL (2019) New taxa and new records of Erythroneurini from China (Hemiptera: Cicadellidae: Typhlocybinae). *Acta Entomologica Musei Nationalis Pragae* 59(1): 189–210. <https://doi.org/10.2478/aemnp-2019-0017>
- Dietrich CH (2005) Keys to the families of Cicadomorpha and subfamilies and tribes of Cicadellidae (Hemiptera: Auchenorrhyncha). *The Florida Entomologist* 88(4): 502–517. [https://doi.org/10.1653/0015-4040\(2005\)88\[502:KTTFOC\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2005)88[502:KTTFOC]2.0.CO;2)
- Dmitriev DA, Anufriev GA, Bartlett CR, Blanco-Rodríguez E, Borodin OI, Cao YH, Deitz LL, Dietrich CH, Dmitrieva MO, El-Sonbati SA, Evangelista de Souza O, Gjonov IV,

- Gonçalves AC, Hendrix S, McKamey S, Kohler M, Kunz G, Malenovský I, Morris BO, Novoselova M, Pinedo-Escatel JA, Rakitov RA, Rothschild MJ, Sanborn AF, Takiya DM, Wallace MS, Zahniser JN (2022) [onward] World Auchenorrhyncha Database. Taxon-Pages. <https://hoppers.speciesfile.org> [Retrieved on 2024-02-21]
- Dworakowska I (1972) On some Oriental Erythroneurini (Auchenorrhyncha, Cicadellidae, Typhlocybinae). Bulletin de l'Académie Polonaise des Sciences. Série des Sciences Biologiques 20(6): 395–405.
- Dworakowska I (1993) Remarks on *Alebra* Fieb. And Eastern Hemisphere Alebrini (Auchenorrhyncha: Cicadellidae: Typhlocybinae). Entomotaxonomia 15(2): 91–121.
- Dworakowska I, Viraktamath CA (1975) On some Typhlocybinae from India (Auchenorrhyncha, Cicadellidae). Bulletin de l'Académie Polonaise des Sciences. Série des Sciences Biologiques 23(8): 521–530.
- Jiang J, Luo GM, Song YH (2021) Two new species of the genus *Arboridia* Zachvatkin from China (Hemiptera: Cicadellidae: Typhlocybinae). Zootaxa 3(3): 349–357. <https://doi.org/10.11646/zootaxa.5005.3.9>
- Matsumura S (1932) A revision of the Palaearctic and Oriental Typhlocybid-genera with descriptions of new species and new genera. Insecta Matsumurana 6(3): 93–120.
- Pu TY, Wang JQ, Song YH (2023) Two new species of Erythroneurini from China (Hemiptera: Cicadellidae: Typhlocybinae). Zootaxa 5374(2): 295–300. <https://doi.org/10.11646/zootaxa.5374.2.8>
- Sohi AS, Sandhu PK (1971) *Arborifera* - a new subgenus of *Arboridia* Zachv. (Typhlocybinae, Cicadellidae) from Punjab, India, with description of its immature stages. Bulletin de l'Académie Polonaise des Sciences. Série des Sciences Biologiques 19(6): 401–406.
- Song YH, Li ZZ (2013) Some new species and new record of the genus *Arboridia* Zachvatkin (Hemiptera: Cicadellidae: Typhlocybinae) from six provinces of China. Zootaxa 3613(3): 229–244. <https://doi.org/10.11646/zootaxa.3613.3.2>
- Song YH, Li ZZ, Dietrich CH (2016) Contribution to knowledge of Typhlocybinae (Hemiptera: Cicadellidae) in Thailand: first record of genus *Arboridia* Zachvatkin with description of three new species and a new synonymy. Zootaxa 4171(2): 357–364. <https://doi.org/10.11646/zootaxa.4171.2.8>
- Zachvatkin AA (1946) Studies on the Homoptera of Turkey. I–VII. Transactions of the Royal Entomological Society of London 97(6): 149–176. <https://doi.org/10.1111/j.1365-2311.1946.tb00278.x>
- Zhang N, Jiang J, Song YH (2022) Two new species of the genus *Arboridia* Zachvatkin from karst area of southwestern China (Hemiptera: Cicadellidae: Typhlocybinae). Journal of Asia-Pacific Entomology 25(3): 101970. <https://doi.org/10.1016/j.aspen.2022.101970>