

Redescription of the enigmatic neotropical inquiline *Paramyrmex foveipennis* Bruch, 1929 with notes on myrmecophily (Coleoptera, Histeridae) in the Sapriniinae subfamily

Tomáš Lackner¹

¹ *Bavarian State Collection of Zoology, Münchhausenstraße 21, 81247 Munich, Germany*

Corresponding author: *Tomáš Lackner* (tomaslackner@me.com)

Academic editor: *M. Caterino* | Received 13 March 2017 | Accepted 25 April 2017 | Published 19 May 2017

<http://zoobank.org/D4AA72BA-B86D-4DFE-940B-252DDB93BCA7>

Citation: Lackner T (2017) Redescription of the enigmatic neotropical inquiline *Paramyrmex foveipennis* Bruch, 1929 with notes on myrmecophily (Coleoptera, Histeridae) in the Sapriniinae subfamily. *ZooKeys* 675: 57–73. <https://doi.org/10.3897/zookeys.675.12690>

Abstract

The poorly-known and highly autapomorphic myrmecophilous Neotropical taxon *Paramyrmex foveipennis* Bruch, 1929 is redescribed, figured and its lectotype designated. Notes on the evolution of the inquiline lifestyle (myrmecophily) in the subfamily Sapriniinae are given.

Keywords

Coleoptera, Histeridae, myrmecophily, *Paramyrmex*, Sapriniinae

Introduction

Several years ago the first higher-level phylogeny of the Sapriniinae was published, where all but three genera and subgenera of this subfamily were included (Lackner 2014). The outstanding taxa were *Auchmosaprinus* Wenzel, 1962, a subgenus of *Xerosaprinus* Wenzel, 1962; *Satrapister* Bickhardt, 1912; and *Paramyrmex* Bruch, 1929. *Satrapister* was the subject of a separate paper (Lackner 2016), whereas *Paramyrmex* is the subject of the present study. The monotypic genus *Paramyrmex* was described by Carlos Bruch based

on five specimens collected by himself and Prof. J. Hubrich in 1923 and 1927, respectively, inside refuse chambers of nests of the ant *Pogonomyrmex cunicularis* var. *carnivora* Santschi, 1925 (= currently a junior synonym of *Pogonomyrmex serpens* Santschi, 1922) in the Santa Fé province of Argentina (Bruch 1929). Bruch had difficulties with the systematic placement of this species, and, referring to the key to histerid genera available to him at the time (Bickhardt 1916), placed the taxon into the Sapriniinae subfamily between ant-inquilines *Myrmetes* Marseul, 1862 and *Platysaprinus* Bickhardt, 1916. The same author noted the peculiarities of *Paramyrmetes*: the rounded, tongue-like labrum and rectangular head, with a broad protruding clypeus that is fused with the frons, and the absent frontal stria. The position of the antennae placed pressed against and parallel to the prosternal process when the head is retracted as well as the prosternum itself resembles, according to Bruch, the taxon *Platysaprinus* (currently a subgenus of *Euspilotus* Lewis, 1907). Bruch also noted several autapomorphies of this genus: the presence of elytral depressions as well as reduced dorsal elytral striae. He pointed out the structure of the antennae and the dilated tibiae as putative morphological adaptations to myrmecophily and observed that *Paramyrmetes foveipennis* was the first recorded (beetle) ant-guest of *Pogonomyrmex cunicularis* var. *carnivora* (= *Pogonomyrmex serpens* Santschi, 1922). In his catalogues (1984, 1997, 2011) Mazur consistently placed *Paramyrmetes* between the Australian *Tomogenius* Marseul, 1862 and Palaeartic *Myrmetes* Marseul, 1862.

In this paper, *Paramyrmetes* is re-described based on the type material. Habitus images as well as drawings of male genitalia are provided. This work represents another contribution to the systematics and higher taxonomy of the Sapriniinae (see e.g. Lackner 2014 or 2016 and the references therein).

Material and methods

A dry-mounted syntype of *Paramyrmetes foveipennis* was relaxed in warm water for several hours. After removal from the original card, it was side-mounted on a triangular point and examined under a Nikon 102 binocular microscope and viewed with diffuse light. Male genitalia were first macerated in 10% KOH solution for about 3 hours, cleared in 80% alcohol and macerated in lactic acid with fuchsine, incubated at 60 °C for another 30 minutes, and subsequently cleared in 80% alcohol and then observed in α -terpineol in a small dish. Digital photographs of male genitalia were taken by a Nikon 4500 Coolpix camera and edited in Adobe Photoshop CS5. Genitalia drawings based on the photographs, or direct observations were produced with the aid of Hakuba klv-7000 light box. Habitus photographs were taken by F. Slamka (Bratislava, Slovakia). The specimen was measured with an ocular micrometer. Beetle terminology follows that of Ôhara (1994) and Lackner (2010).

The specimen examined for this study is deposited in the following collection:

MNNC Museo Nacional de Historia Natural (Santiago de Chile, Chile).

Abbreviations. Abbreviations of morphological measurements follow Ôhara (1994) and are used throughout the text as follows:

APW	width between anterior angles of pronotum
EL	length of elytron along elytral suture
EW	maximum width between outer margins of elytra
PEL	length between anterior angles of pronotum and apices of elytra
PPW	width between posterior angles of pronotum.

Results

Paramyrmex Bruch, 1929

Paramyrmex Bruch, 1929: 421. Type species *Paramyrmex foveipennis* Bruch, 1929: 422, by monotypy.

Paramyrmex: Mazur (1984): 107; Mazur (1997): 217; Mazur (2011): 178.

Diagnosis. Medium-sized reddish-brown shining asetose Sapriniinae beetle with completely punctate and shagreened dorsal cuticle, broadly rectangular head; frontal and supraorbital striae absent, labrum tongue-shaped. Dorsal elytral striae strongly reduced; apical third of elytra deeply depressed; metaventrite and first visible abdominal ventrite with (striolate) depression. Pygidium with prominent round ornamentation; tibiae dilated.

Differential diagnosis. Based on the autapomorphies outlined above, *Paramyrmex* cannot be confused with any currently known South American Sapriniinae genus. The overall body coloration, in combination with the depressed apical third of the elytra, metaventral and abdominal depressions and, especially the peculiarly-shaped labrum, will readily set this taxon apart from the other members of the subfamily. Moreover, *Paramyrmex* possesses pygidial ornamentation in the male sex. Pygidial ornamentation occurs in the Sapriniinae subfamily rather seldom, and was observed so far only with female specimens of several taxa (e.g. *Euspilotus* (*Neosaprinus*) *perrisi* Marseul, 1872)). According to my knowledge, male pygidial ornamentation has not been reported in the Sapriniinae subfamily hitherto.

Biology. The type series was found inside the refuse chambers of the ant *Pogonomyrmex serpens* Santchi, 1922. This species is apparently a specialised ant inquiline.

Distribution. Known only from the type series collected in the province of Santa Fé, Argentina (Fig. 17).

Remarks. Although Bruch (1929) mused about the metaventral- and abdominal depressions as a possible sexual character of *Paramyrmex*, he did not specify the sex of the specimens he examined. As I was only able to examine a single male, I am not sure whether *Paramyrmex foveipennis* is a sexually dimorphic species or not.

***Paramyrmex foveipennis* Bruch, 1929**

Figs 1–17

Paramyrmex foveipennis Bruch, 1929: 422, figs 1–5, 11; Mazur (1984): 107; Mazur (1997): 217; Mazur (2011): 178.

Type material examined. Lectotype, present designation, ♂, side-mounted on a triangular mounting card, genitalia dismembered, glued to a separate card under the specimen, with the following labels: “Hersilia / “La Geraldina”” (written); followed by: “Prov.S.Fé.II.927 / J. Hubrich” (written); followed by: “con / Pogonomymex / v.carnivora” (written); followed by: “Phototypus” (light-green label, written); followed by: “Paramyrmex / foveipennis / Bruch / C. BRUCH DETERM.” (printed-written); followed by: “ACHADO” (written); followed by: “Paramyrmex / foveipennis / Bruch, 1929 / LECTOTYPE / Des. T. Lackner 2017” (red label, written) (MNNC). This specimen is undoubtedly one of the syntypes, as Bruch described the species based on a single specimen collected by him and four specimens collected by Prof. Hubrich in Hersilia. The examined specimen belongs to the series collected by Prof. Hubrich. According to Bruch (1929) the type specimens were deposited in his private collection, and the private collections of Reichensperger and Hubrich. It is possible that the private collection of Prof. Hubrich was acquired by MNNC, where the lectotype is housed currently. Lectotype designation fixes the species’ identity; the outstanding four exemplars, whose whereabouts are unknown, would qualify for the paralectotype status.

Re-description. Body: PEL: 3.00 mm; PPW: 1.00 mm; APW: 2.00 mm; EW: 2.60 mm; EL: 1.70 mm. Body, including appendages reddish-brown (Figs 1–2). Head broad, frons approximately twice as wide as long, with several vague depressions, frontal and supraorbital striae absent; clypeus large, quadrate, sloping down laterally; both frons and clypeus covered with dense fine punctures separated by approximately 1–2 times their diameter, interspaces with alutaceous microsculpture. Eyes flattened, but visible from above; antennal scape slender, with several setae; pedicel approximately as long as following three antennomeres together; 8th antennomere saucer-like. Antennal club oval, on distal half (roughly) with dense microsetae intermingled with several longer setae; sensory structures not examined. Mouthparts: labrum (Fig. 3) unusually large, tongue-shaped; labral pits situated on labral edges, each with two labral setae. Mandibles similar to other taxa of the subfamily, apically pointed, finely punctate. Ultimate palpomere of both labial and maxillary palpi elongate, thin, approximately four times as long as wide. Mentum quadrate, anterior margin with deep semi-circular median emargination; rest of mouthparts not examined. Pronotum broad, approximately twice as wide as long; marginal pronotal striae carinate laterally, weakened behind head. Apical pronotal angles obtuse; pronotal disk with punctation similar to that of head, punctures fine, separated by approximately twice their diameter, interspaces with prominent alutaceous microsculpture. Pronotal hypomeron glabrous. Elytral epipleuron with microscopic punctation, marginal epi-




Figure 1. *Paramyrmex foveipennis* Bruch, 1929, ♂, lectotype, habitus, dorsal view.

pleural stria very fine, complete; marginal elytral stria complete, carinate, continued as weakened but complete marginal elytral stria connected to fine, complete sutural elytral stria. Humeral elytral stria fine, doubled, present on basal elytral third approximately; both subhumeral striae lacking. First dorsal elytral stria shortened on approximate basal tenth and apical fifth; second dorsal elytral stria even shorter, shortened on approximate basal fifth and apical third; rest of the elytral striae (apart from fine and complete sutural elytral stria) missing. Elytral disk in anterior two-thirds relatively coarsely, subimbricately punctate, punctures with posterior margins effaced, each with short appressed seta one to two times puncture diameter, interpunctural integument with very dense and prominent alutaceous microsculpture; in posterior third with punctation similar to that of pronotum and near elytral suture with rather deep large circular depression (Fig. 4). Scutellum very small, triangular. Propygidium partly covered by elytra, densely punctate, punctures separated by about their own



Figure 2. *Paramyrmex foveipennis* Bruch, 1929, ♂, ditto, ventral view.

diameter; pygidium triangular, densely punctate, punctures finer than those of propygidium, separated about their own diameter; apex of pygidium (Fig. 5) with round prominent well-defined depression, depression edges roughly subcarinate, with raised narrow triangular inclusion beginning at basomedial third of margin and ending at apical fifth, inclusion subcarinate in its apical two-thirds.




Figure 3. *Paramyrmex foveipennis* Bruch, 1929, ♂, head, dorsal view, showing the peculiar labrum (redrawn from Bruch 1929).



Figure 4. *Paramyrmex foveipennis* Bruch, 1929, ♂, lectotype, detail of the elytral depression.



Figure 5. *Paramyrmex foveipennis* Bruch, 1929, ♂, ditto, pygidium, caudal view.



Figure 6. *Paramyrmex foveipennis* Bruch, 1929, ♂, ditto, prosternum and mesoventrite.





Figure 7. *Paramyrmex foveipennis* Bruch, 1929, ♂, ditto, schematic detail of the prosternal process showing the curious circle presumed to help with the accommodation of antennal club when head is retracted.

Prosternum: apical margin of prosternum (Fig. 6) rounded, marginal prosternal stria absent; carinal prosternal striae present, largely divergent apically; surface between striae convex. Lateral prosternal striae costate, apically fusing with widely divergent apices of carinal prosternal striae slightly anterior to anterior third; prosternal foveae absent. Prosternal process laterally with a single large dark circular depression presumed to accommodate antennal club when head is retracted (Fig. 7). Mesoventrite: lateral mesoventral stria complete, inwardly arcuate medially. Punctuation of mesoventral disk similar to that of pronotum, punctures separated by 1-2 times their diameter, interspaces with microsculpture. Metaventrte: sparsely and finely punctate, lateral metaventral stria absent. Basal half of metaventral disk moderately depressed, this depression continues and occupies entire first visible abdominal ventrite which is strongly striolate (Fig. 2). Lateral disk of metaventrte with subcontiguous large shallow punctures, metepisternum with similar punctuation; metepisternal stria thin, complete.

Legs: protibia dilated, outer margin with approximately 14 short denticles, teeth lacking. Protibial spur thin, emerging near tarsal insertion, protarsal groove shallow. Mesotibia and metatibia not particularly dilated, outer margin with sparse denticles.

Male genitalia: 8th sternite (Figs 8–9) apically without setae; 8th tergite (Fig. 9) apically outwardly arcuate; 8th sternite and tergite laterally fused (Fig. 11). Tenth tergite apically inwardly arcuate (Fig. 12); 9th tergite fused dorsally (Fig. 12). Spiculum gastrale (9th sternite) gradually dilated on both ends (Figs 13–14). Aedeagus (Figs 15–16) curved in lateral view, phallobase approximately one fifth of tegmen's length. Parameres fused on their basal two-thirds. Female unavailable.



Figures 8–16. 8 *Paramyrmex foveipennis* Bruch, 1929, ♂, lectotype, terminalia: eighth sternite and tergite, ventral view 9 ditto, dorsal view 10 ninth and tenth tergites, lateral view 11 eighth sternite and tergite, lateral view 12 ninth and tenth tergites, dorsal view 13 spiculum gastrale (ninth sternite), ventral view 14 ditto, lateral view 15 aedeagus, dorsal view 16 ditto, lateral view.



Figure 17. Distribution of *Paramyrmetes foveipennis* Bruch, 1929 in Argentina.

Discussion

In his catalogues Mazur (1984, 1997, 2011), without any prior analysis, consistently placed *Paramyrmetes* among the more ‘basal’ Sapriniinae, between the cave-dwelling and inquilinous Australopacific genus *Tomogenius* Marseul, 1862 and Palaearctic ant-inquiline genus *Myrmetes* Marseul, 1862. In my analysis, both *Tomogenius* and *Myrmetes* were found to be near the root of the tree, and, in fact an inquilinous lifestyle of saprinines is inferred to be plesiomorphic for the subfamily (Lackner 2014).

For the present study, I coded the morphological characters of *Paramyrmetes* and included them in the matrix used for my *Satrapister* paper (Lackner 2016). The position of *Paramyrmetes* on the tree was unstable, varying among the taxa near the root (*Myrmetes*, *Erebidus*, *Gnathoncus*, *Tomogenius*, *Microsaprinus*). This instability was mainly due to the impossibility of coding antennal club characters, as the disarticulation of the unique syntype available was not allowed. Furthermore, *Paramyrmetes* exhibits numerous autapomorphies, which are not informative for the phylogenetic reconstruction. Consequently, I decided not to include the new tree here, opting instead to await the results of the molecular analysis (on-going). Despite the unstable position on the tree, *Paramyrmetes*, consistently came out near the root of the tree, in fact corroborating Mazur’s taxon placement. Inclusion of freshly collected specimens into the ongoing molecular analysis would be highly desirable.


Myrmecophily, or an ant-inquilinous lifestyle is rather rare among saprinine histerids and has hitherto been confirmed in six higher taxa with at least three independent evolutionary events leading to it (Lackner 2014; P. Kovarik, pers. communication). It is interesting to note that four out of the six ant-inquiline higher taxa occur in the New World, and three of those are attaphilic (see the table below for details). Several, but not all inquilinous saprinines exhibit peculiar morphological autapomorphies linked to their specialized lifestyles, and these autapomorphies are not similar between the taxa. Among the shared characters could be listed their wholly punctate bodies (except for *Myrmetes*), interrupted to absent frontal stria (all taxa), and occasional lack of sutural elytral stria (present in *Priscosaprinus* and *Paramyrmetes*). The dilated tibiae, which are present in most taxa (exceptions are: *Myrmetes*, *Priscosaprinus* and *Phoxonotus*) probably function to protect larger portion of venter when in repose, since they are able to cover more space. Autapomorphies of the undescribed Australian genus, which include imbricately setose elytra and abdomen (it is hypothesized that these micro-setae function as trichomes secreting appeasing liquid), metepisternal groove for receiving the mesotarsus, and enlarged antennal club are further results of the selection pressures for inquilinous lifestyle. Many of these are also found in other ant-inquiline histerids (e.g. Haeteriinae). The functions of several autapomorphies, e.g. elytral and metepisternal-abdominal depressions, dorsal tubercles, tongue-shaped labrum or pygidial ornamentation is completely unknown. Larvae of ant-inquiline saprinine beetles are unknown and biology generally poorly understood. The table below is presented to outline several putative morphological adaptations presumably linked with ant-inquilinism in the Sapriniinae subfamily; a dorsal habitus image of a representative of each taxon is included.

Taxon name	No of ant-inquilineous species	Ant genus or species association	Distribution	Morphological adaptations (linked to ant-inquility?)
<i>Myrmetes</i> Marseul, 1862 (Fig. 18)	1	<i>Formica</i> sp.	Western Palaearctic	Absent sutural stria; absent protibial groove; absent frontal stria; body impunctate.
New genus (Fig. 19)	1	<i>Iridomyrmex purpureus</i> (Smith, 1858)	Australia: New South Wales, Queensland	Absent inner subhumeral and sutural stria; frontal stria prolonged onto clypeus; elytra imbricate with microscopic setae; 5 carinate elytral striae; abdominal ventrites and sternites micro-setose; tibiae dilated; metepisternum with groove for tarsus; wholly punctate body.
<i>Paramyrmetes</i> Bruch, 1929 (Fig. 1)	1	<i>Pogonomyrmex serpens</i> Santschi, 1922	Argentina	Absent frontal stria; tongue-shaped labrum; elytral depression, reduced elytral striation; pygidial ornamentation; dilated tibiae; metaventrite & first abdominal ventrite depressed; wholly punctate body.
<i>Phoxonotus</i> Marseul, 1862 (Fig. 20)	5	<i>Atta sexdens</i> Linnaeus, 1758; <i>A. cephalotes</i> Linnaeus, 1758	Guatemala, Costa Rica, French Guyana, Suriname, Peru, Brazil	Tubercles on dorsum, deeply incised prosternal base; absent sutural stria; frontal stria interrupted, prolonged onto clypeus; wholly punctate body.
<i>Euspilotus (Platysaprinus)</i> Bickhardt, 1916 (Fig. 21)	2	<i>Acromyrmex</i> sp. (Antini)	Argentina, Brazil	Frontal stria interrupted, prolonged onto clypeus; longitudinal tubercle between elytral suture and 4 th stria; absent/present sutural stria; dilated tibiae; wholly punctate body.
<i>Geomyssaprinus (Priscosaprinus)</i> Wenzel, 1862 (Fig. 22)	1	<i>Atta texana</i> (Buckley, 1860)	Mexico, USA: Louisiana	Frontal stria interrupted; otherwise no particular autapomorphies.



18

Figure 18. *Myrmetes paykulli* Kanaar, 1979 habitus, dorsal view.



19

Figure 19. Undescribed genus, Australia habitus, dorsal view.



Figure 20. *Phoxonotus (Ph.) parvotuberculatus* Lackner, 2016 habitus, dorsal view.



Figure 21. *Euspilotus (Platysaprinus) latimanus* (Schmidt, 1890) habitus, dorsal view.



Figure 22. *Geomysaprinus (Priscosaprinus) formicus* (Hinton, 1935) habitus, dorsal view.

Acknowledgements

I thank Gerardo Arriagada (Santiago de Chile, Chile) for his help acquiring the lectotype of *Paramyrmex foveipennis* for my study. Funding for this study has been provided by the Alexander von Humboldt Foundation (Bonn, Germany), itself part of the Federal Ministry for Education and Research (Berlin, Germany). I thank Alexey K. Tishechkin (Washington D.C., USA), Peter W. Kovarik (Columbus, USA), William B. Warner (Chandler, USA) as well as the editor for the Histeroidea at ZooKeys for their critical comments and grammar correction of the manuscript. I thank my wife Pepina for re-drawing the figure from Bruch (1929) in Adobe Illustrator. Alexey K. Tishechkin is being thanked for providing a photograph of *Geomysaprinus (Priscosaprinus) formicus* used in this paper.

References

- Bickhardt H (1916) Histeridae. In: Wytzman P (Ed.) *Genera Insectorum*, Fascicle 166a, b. La Haye, 113–302.
- Bruch C (1929) Neue myrmekophile Histeriden und Verzeichnis der aus Argentinien bekannten Ameisengäste. *Zoologischer Anzeiger* 82: 421–437.
- Lackner T (2014) Review of the Palearctic genera of Sapriniinae (Coleoptera: Histeridae). *Acta Entomologica Musei Nationalis Pragae* 50(Supplementum): 1–254.

- Lackner T (2014) Phylogeny of the Sapriniinae subfamily reveals interesting ecological shifts in the history of the subfamily (Coleoptera: Histeridae). *Zoological Journal of the Linnean Society* 172: 521–555. <https://doi.org/10.1111/zoj.12182>
- Lackner T (2016) *Satrapister nitens* Bickhardt, 1912: redescription and tentative phylogenetic placement of a mysterious taxon (Coleoptera, Histeridae, Sapriniinae). *Deutsche Entomologische Zeitschrift* 63(1): 1–8. <https://doi.org/10.3897/dez.63.6363>
- Mazur S (1984) A world catalogue of Histeridae. *Polskie Pismo Entomologiczne* 54(3-4): 1–376.
- Mazur S (1997) A world catalogue of the Histeridae. *Genus – International Journal of Invertebrate Taxonomy (supplement)*, 373 pp.
- Mazur S (2011) A concise catalogue of the Histeridae. Warsaw University of Life Sciences, SGGW Press, Warsaw, 332 pp.
- Ôhara M (1994) A revision of the superfamily Histeroidea of Japan (Coleoptera). *Insecta Matsumurana (NS)* 51: 1–283.