

A new species of *Atractides* Koch, 1837 (Acari, Hydrachnidia, Hygrobatidae) from Ethiopia, with a discussion on the biodiversity of the genus *Atractides* in the Afrotropical region

Vladimir Pešić^{1,†}, Harry Smit^{2,‡}

1 Department of Biology, University of Montenegro, Cetinjski put b.b., 81000 Podgorica, Serbia and Montenegro **2** Zoological Museum, University of Amsterdam, Plantage Middenlaan 64, 1018 DH Amsterdam, The Netherlands

† <urn:lsid:zoobank.org:author:719843C2-B25C-4F8B-A063-946F53CB6327>

‡ <urn:lsid:zoobank.org:author:A610019D-74C5-441C-8F01-E39A03622EAE>

Corresponding author: Vladimir Pešić (pesicv@t-com.me)

Academic editor: Andre Bochkov | Received 27 January 2011 | Accepted 10 March 2011 | Published 19 March 2011

<urn:lsid:zoobank.org:pub:9BE15F61-7256-4E91-9E34-BC464C9C8296>

Citation: Pešić V, Smit H (2011) A new species of *Atractides* Koch, 1837 (Acari, Hydrachnidia, Hygrobatidae) from Ethiopia, with a discussion on the biodiversity of the genus *Atractides* in the Afrotropical region. ZooKeys 86: 1–10. doi: [10.3897/zookeys.86.972](https://doi.org/10.3897/zookeys.86.972)

Abstract

A new species of *Atractides* Koch, 1837 (Acari, Hydrachnidia) is described from Ethiopia. The world number of *Atractides* now tallies 297 species. The diversity of the genus *Atractides* in the Afrotropical region is briefly discussed.

Keywords

Water mites, taxonomy, new species, Africa, biodiversity

Introduction

Water mites of the genus *Atractides* Koch, 1837 have been found in all biogeographical regions except Australia and Antarctica. Gerecke (2003) reported 123 species from Europe, 72 from Asia, 27 from North America, 33 from Central and South America

and 54 from Africa. Since then, many new species have been described (Pešić et al. 2004, Pešić and Smit 2009, Tuzovskij 2011), suggesting that many more remain to be discovered.

The aim of our paper is to describe a new *Atractides* species from Ethiopia. Additionally, the worldwide biodiversity of the genus *Atractides* Koch, 1837 of the world, with an emphasis on the Afrotropical region, is briefly discussed.

Material and methods

Water mites were collected by hand netting, sorted on the spot from the living material, preserved in Koenike fluid and dissected as described elsewhere (e.g. Davids et al. 2007). Holotype and paratypes are deposited in the Zoological Museum in Amsterdam (ZMAN). All material has been collected by the junior author.

All measurements are given in μm . For a detailed description and discussion of the characteristics of the genus *Atractides* and a detailed methodological introduction, see Gerecke (2003).

The following abbreviations are used: Ac-1 = first acetabulum, alt. = altitude, asl. = above sea level, Cx-I = first coxae, dL = dorsal length, H = height, HB = central height, L = length, I-L-6 = Leg 1, sixth segment (tarsus), lL = lateral length, mL = medial length, P-1 = palp, first segment, S-1 = large proximal ventral seta at I-L-5, S-2 = large distal ventral seta at I-L-5, Vgl = ventroglandulare, vL = ventral length, W = width.

Systematics

Atractides (Atractides) ethiopiensis sp. n.

urn:lsid:zoobank.org:act:E968E96B-BAF1-4641-8DB4-3490FC07C28C

Figs 1–3

Type series. Holotype, male (ZMAN), dissected and slide mounted, Ethiopia, Roby River, 21.x.2006, $9^{\circ}44.996\text{N}$, $38^{\circ}59.743\text{E}$, 2507 m a.s.l. Paratypes: two males, one female (ZMAN, one female dissected and slide mounted), same data as holotype.

Diagnosis. Dorsal integument striated, palp slender with P-2 straight in the both sexes, S-1 in female ending in a fine hair-like tip, excretory pore surrounded by distinct oval sclerite, Vgl-1 fused to Vgl-2.

Description. General features. Dorsal integument: striated, muscle attachment plates unsclerotized. Coxal field: extended secondary sclerotization, caudal margin Cx-I broadly convex. Genital field: Ac in a weakly curved line. Excretory pore: sclerotized; Vgl-1 fused to Vgl-2. Palp: ventral margin of P-2 and ventral margin of P-3 slightly concave, ventral margin of P-4 straight, sword seta between ventral hairs, nearer to the distoventral hairs. Legs: I-L-5 S-1 and S-2 pointed, interspaced, S-2 basally enlarged,




Figure 1. *Atractides ethiopiensis* sp. n., male: idiosoma, ventral view. Scale bar = 100 μm .

bluntly pointed; I-L-6 slender, curved, with maximum H proximally; leg claws with dorsal and ventral clawlets (Fig. 2C).

Morphology. Male: Idiosoma L/W 669-684/541-556; coxal field L 420; Cx-III W 484; Cx-I+II mL 156; IL Cx-I+II 265. Genital field (Fig. 1): anterior margin convex, with a border of secondary sclerotization convexely protruding, anterior margins of gonopore and Ac-1 considerably distant from anterior margin of genital field; L/W 150/150; L Ac 1-3: 38-41, 39-45, 39-41.

Palp (Fig. 2A): total L 369, dL: P-1, 35; P-2, 72; P-3, 92; P-4, 131; P-5, 39; L ratio P-2/P-4, 0.55; P-4 club-shaped, with maximum H near distoventral hair.

I-L: I-L-5 dL 243, vL 169, dL/vL ratio 1.44, HB 50, dL/HB 4.86, S-1 L 121, L/W 11.2, S-2 L 95, L/W 5.6, distance S-1-2 23, L ratio S-1/2 1.27; I-L-6 L 166, HB 16, L/HB ratio 10.3; L ratio I-L-5/6 1.46.

Female. Idiosoma L/W 1106/928; coxal field L 473; Cx-III W 644; Cx-I+II mL 147; IL Cx-I+II 284. Genital field (Fig. 3B): L/W 181/209; genital plate bean-shaped, with slightly concave medial margins, L 139; L Ac 1–3: 44, 47, 43.

Palp (Fig. 2D): total L 417, dL: P-1, 48; P-2, 80; P-3, 115; P-4, 132; P-5, 42; L ratio P-2/P-4, 0.6; P-4 not club-shaped, with maximum H near proximoventral hair; chelicera total L 250, claw L 83.

I-L: I-L-5 dL 279, vL 186, dL/vL ratio 1.5, HB 65, dL/HB 4.3, S-1 ending in a fine hair-like tip (Fig. 3C) L 162, L/W 11.3, S-2 L 117, L/W 4.8, distance S-1-2 29, L ratio S-1/2 1.38; I-L-6 L 210, HB 19, L/HB ratio 11.1; L ratio I-L-5/6 1.33.

Etymology. Named after its occurrence in Ethiopia.

Remarks. Due to the striated integument, a slender palp with P-2 lacking ventro-distal projection and similar morphology of I-L-5 and -6, with S-1 in female ending in a fine hair-like tip *Atractides ethiopiensis* sp. n. resembles *A. latisetus* (K. Viets, 1916), a species known from Cameroon (K. Viets 1916), Liberia (Cook 1966), East and South Africa (Lundblad 1952, Viets 1964). This species can be easily distinguished from *Atractides ethiopiensis* sp. n., by a smooth excretory pore and unfused Vgl-1 and -2.

Distribution. Ethiopia.

Discussion

a) The present biodiversity of *Atractides*

To get an overview of the biodiversity of the genus worldwide, we examined numerous papers, Gerecke (2003) and the website www.watermite.org (viewed on January 23, 2011). The total number of *Atractides* species worldwide tallies exactly 297 species, including the new species described in this paper. Most species are known from the northern Hemisphere: 138 are described from the Palaearctic, most of these from the Western Palaearctic and the countries surrounding the Mediterranean Sea (Gerecke 2003). In the Oriental region 56 species have been found, 47 species have been found in the Afrotropical region, including the new species from Ethiopia, while 32 and 27 species are described from the Neotropical and Nearctic region, respectively (Fig. 4).

b) The biodiversity of *Atractides* in Afrotropical region

Thus far, 47 species and 3 subspecies have been recorded from the Afrotropical region, with large differences in the state of knowledge of different regions. The number of known species ranges from 14 from Kenya, 13 from South Africa and 9 from Liberia (Viets 1970; Jansen van Rensburg 1974). Conversely, only two species, *Atractides con-*




Figure 2. *Atractides ethiopiensis* sp. n., (A–C male, D female) **A, D** palp, medial view **B** I-L-5 and -6 **C** leg claw. Scale bar = 100 µm.

temptus (Lundblad, 1951) and *A. ethiopiensis* sp. n. are known from Ethiopia (Motaš and Tanasachi 1968, present paper).

The Afrotropical members of *Atractides* belong to the following subgenera *Atractides* Koch, 1837 s.s., *Megabates* K. Viets, 1924, *Tympanomegapus* Thor, 1923 and *Polymegapus* K. Viets, 1926. However, the older data (Viets 1970; Jansen van Rensburg 1974), as well as the most recent catalogue of water mites (Viets 1987) assigned *Tympanomegapus* and triacetabulate members of *Polymegapus* to *Atractides* Koch, 1837 s.s.

A critical analysis of the Afrotropical *Atractides* species, with the aid of the diagnoses and the revised key presented in Gerecke (2003), shows that nine species (i.e., *Atractides harrisoni* K.O. Viets, 1971, *A. levipapis* Bader, 1968, *A. neotestudo* Cook, 1966, *A. paratestudo* Cook, 1966, *A. pseudotestudo* Cook, 1966, *A. scutifer* (Lundblad, 1951), *A. subtestudo* Cook, 1966, *A. testudo* Cook, 1966 and *A. tuberipalpis* (K. Viets 1913)) should be assigned to *Tympanomegapus*, while four species (i.e., *A. abruptus* Cook, 1966, *A. congoensis* K.O. Viets & Böttger, 1972, *A. kuhnei* (K. Viets, 1911) and *A. multiporus* Cook, 1966) should be assigned to the subgenus *Polymegapus*. The subgeneric position of *Atractides laetus* K.O. Viets & Böttger, 1972 is unclear. Due to P-1 by far longer than high and centrally narrowed, this species agrees with members of *Tympanomegapus*, but differs in a stout cheliceral basal segment (L/H 2.4, calculated from Viets and Böttger 1972) and a remarkably long claw (Viets and Böttger 1972).

The subgenus *Megabates* K. Viets, 1924, includes two Afrotropical species, i.e., *Atractides rectipes* (K. Viets, 1924) and *A. longicoxalis* (Cook, 1974). According to Gerecke (2003), *Megabates* is most probably a synonym of *Atractides*.

The other 35 species and subspecies are assigned to *Atractides* s. s. According to our present state of knowledge, slightly more than a half (54%) of these species are known from both sexes, i.e., *Atractides comorosensis* Smit & Pešić, 2010, *A. contemptus* (Lundblad, 1951), *A. coriacellus* K. Viets, 1956, *A. damkoheleri* (K. Viets, 1916), *A. ethiopiensis* sp. n., *A. falcipes* (Walter & Bader, 1952), *A. kilimandjaricus* Lundblad, 1952, *A. latisetus* (K. Viets, 1916), *A. linearis* (Lundblad, 1927), *A. lundbladi lundbladi* (Halik, 1947), *A. madagascariensis* K.O. Viets, 1964, *A. minutissimus* (Lundblad, 1927), *A. processiferus* (Walter & Bader, 1952), *A. pusillus* (Walter & Bader, 1952), *A. scutelliferus* K.O. Viets, 1964, *A. splendidus splendidus* (Lundblad, 1927), *A. splendidus superbus* (Lundblad, 1927), *A. thoracatus* Koenike, 1898 and *A. valididens* (Lundblad, 1951). Three species are known from the male only, i.e. *Atractides africanus* (Lundblad, 1951), *A. baderi* K. Viets, 1956 and *A. invidendus* K.O. Viets, 1964, while 12 species are known from the female only, i.e., *A. assimilis* K.O. Viets, 1964, *A. callosus* K.O. Viets, 1972, *A. exiguum* Lundblad, 1952, *A. immodestus* (Walter & Bader, 1952), *A. irangiensis* K.O. Viets & Böttger, 1972, *A. kuhlmanni* K.O. Viets, 1963, *A. lundbladi curvitarsis* K. Viets, 1955, *A. pulcher* K. Viets, 1956, *A. rostellatus* K.O. Viets, 1964, *A. sudaficanus* K. Viets, 1956, *A. tenuipes tenuipes* Lundblad, 1952 and *A. tenuipes ambiguus* K.O. Viets, 1971.




Figure 3. *Atractides ethiopiensis* sp. n., female **A** = coxal field **B** = genital field, excretory pore and Vgl-1 and -2 **C** = I-L-5 and -6. Scale bar = 100 μm .




Figure 4. Distribution of freshwater mite species of *Atractides* per biogeographical provinces: PA—Palaeartic; NA—Nearctic; NT—Neotropical; AT—Afrotropical; OL—Oriental; AU—Australasian; PC—Pacific Oceanic Islands; AN—Antarctic. Biogeographical provinces are given according to Balian et al. (2008), modified to include northern Oman in the Oriental, and southern Oman in the Afrotropical region (see Smit and Pešić 2010 for discussion on the zoogeographical position of Oman).

Three species are of doubtful status and merit attention during future taxonomical studies: *Atractides africanus* (Lundblad, 1951) – possibly a synonym of *A. linearis* (Lundblad, 1927) (see Viets 1964 for a discussion); *A. pusillus* (Walter & Bader, 1952) - rather similar to and probably a synonym of *A. damkoehleri* (K. Viets, 1916) (see Viets 1964 for a discussion), and *A. processiferus* (Walter & Bader, 1952) – there is good reason to assume that this species will be found to be synonymous with *A. valididens* (Lundblad, 1951), a species overlooked in the original description of *A. processiferus* (see: Walter and Bader 1952).

In conclusion the current knowledge of the diversity of Afrotropical *Atractides* species is far from complete. Moreover, information on the diversity of Afrotropical *Atractides* among different freshwater habitats is unbalanced, and some important habitats are poorly (e.g., springs) or completely unexplored (e.g., hyporheic interstitial). Additional field work is highly needed for an appropriate evaluation of the extant diversity.

Acknowledgements

We are indebted to Johannes Postma (Ann Arbor) for reviewing the English.

References

- Balian EV, Segers H, Lévèque C, Martens K (2008) An introduction to the freshwater animal diversity assessment (FADA) project. *Hydrobiologia* 595: 3–8.
- Cook DR (1966) The water mites of Liberia. *Memoirs of the American Entomological Institute* 6: 1–418.
- Davids C, Di Sabatino A, Gerecke R, Gledhill T, Van der Hammen H, Smit H (2007) *Acari: Hydrachnidia*. In: Gerecke R (Ed) 2006. Chelicerata: Araraneae, Acari I. Süßwasserfauna von Mitteleuropa 7/2–1: 241–376. Elsevier Spektrum Akademischer Verlag, München.
- Gerecke R (2003) Water mites of the genus *Atractides* Koch, 1837 (Acari: Parasitengona: Hygrobatiidae) in the western Palaearctic region: A revision. *Zoological Journal of the Linnean Society* 138: 141–378.
- Jansen van Rensburg CA (1974) A checklist of the Ethiopian water mites with notes on their distribution. *Wetenskaplike bydraes van die Potchefstroomse Universiteit vir Christelike Hoër Onderwys, Reeks B: Natuurwetenskappe*, 68: 1–51.
- Lundblad O (1951) Vorläufige Beschreibung einiger ostafrikanischer Hydracarinen. *Entomologisk Tidskrift* 72: 157–161.
- Lundblad O (1952) Hydracarinen von den ostafrikanischen Gebirgen. *Arkiv för Zoologi, Serie 2, 3*: 391–525.
- Motaş C, Tanasachi J (1968) Quelques Hydrachnellae (Acari) du Nil Bleu, de la rivière Takaze et du Lac Tana (Ethiopie). *Israel Journal of Zoology* 16: 171–192.
- Pešić V, Saboori A, Asadi M, Vafaei R (2004). Studies on water mites of the family Hygrobatiidae (Acari, Hydrachnidia) from Iran, I. The water mite genus *Atractides* Koch, with the description of five new species. *Zootaxa* 495: 1–40.
- Pešić V, Smit H (2009) New records of water mites of the genus *Atractides* Koch, 1837 (Acari: Hydrachnidia, Hygrobatiidae) from Thailand, Malaysia and Sulawesi (Indonesia), with the description of four new species. *Zootaxa* 2240: 1–30.
- Smit H, Pešić V (2010) New species of water mites from Oman, with some zoogeographical notes (Acari: Hydrachnidia). *Acarologia* 50: 151–195.
- Tuzovskij PV (2011) Four new water mite species of the genus *Atractides* Koch (Acari: Hydrachnidia: Hygrobatiidae) from Russia. *Zootaxa* 2738: 41–59.
- Viets K (1916) Ergänzungen zur Hydracarinen-Fauna von Kamerun (Neue Sammlungen). *Archiv für Hydrobiologie* 11: 241–306.
- Viets K (1956) Wassermilben aus Südafrika. Systematische, faunistische und ökologische Untersuchungen. *Zoologischer Jahrbücher, Abteilung für Systematik, Ökologie und Geographie der Tiere* 84: 1–31.
- Viets KO (1963) Hygrobatiidae (Acari, Hydrachnellae) aus Südafrika. *Archiv für Hydrobiologie* 59: 467–485.
- Viets KO (1964) Neufunde und Taxonomie afrikanischer Hydrachnellae (Acari). *Acarologia* 6: 129–162.

- Viets KO (1970) Unser Zuwachs an Kenntnissen über die aus Afrika bekannten Wassermilben (Hydrachnellae, Acari) (mit Anhang: Limnohalacaridae). *Hydrobiologia* 35: 65–126.
- Viets KO (1971) Wassermilben aus Südafrika (Hydrachnellae, Acari). *Acarologia* 12: 752–779.
- Walter C, Bader C (1952) Hydracarina. In: Mission scientifique de l'Omo. Mémoires du Muséum d'Histoire naturelle, Série A, Zoologie 4: 87–236.